



## Metabolomics of UC bacterial ecosystem compared to the healthy donors

Sulek, Karolina; Vignæs, Louise Kristine; Abbele, P. van den; Frandsen, Henrik Lauritz; Smedsgaard, Jørn; Skov, T. H.; Steenholdt, C.; Brynskov, J.; Wiele, T. van de; Licht, Tine Rask

*Publication date:*  
2012

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Sulek, K., Vignæs, L. K., Abbele, P. V. D., Frandsen, H. L., Smedsgaard, J., Skov, T. H., Steenholdt, C., Brynskov, J., Wiele, T. V. D., & Licht, T. R. (2012). *Metabolomics of UC bacterial ecosystem compared to the healthy donors*. Abstract from 7th Danish Conference on Biotechnology and Molecular Biology, Vejle, Denmark.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Metabolomics of UC bacterial ecosystem compared to the healthy donors.

K. Sulek<sup>1\*</sup>, L. K. Vigsnaes<sup>1</sup>, P. van den Abbeele<sup>2</sup>, H. L. Frandsen<sup>1</sup>, J. Smedsgaard<sup>1</sup>, T. H. Skov<sup>3</sup>, C. Steenholdt<sup>4</sup>, J. Brynskov<sup>4</sup>, T. van de Wiele<sup>2</sup> and T. R. Licht<sup>1</sup>.



<sup>1</sup> DTU Food, Technical University of Denmark, Søborg, Denmark.

<sup>2</sup> Laboratory of Microbial Ecology and Technology (LabMET), Ghent University, Gent, Belgium.

<sup>3</sup> Division of Quality and Technology, Faculty of Life Sciences, University of Copenhagen, Frederiksberg, Denmark.

<sup>4</sup> Department of Medical Gastroenterology, Herlev Hospital, Herlev, Denmark.

\* kasul@food.dtu.dk

## Introduction

Ulcerative colitis<sup>1)</sup> (UC) is an idiopathic inflammatory bowel disease (IBD), which is characterized by chronic inflammation of the colonic mucosa. As the etiology of IBDs remains still unknown, it has been shown in many studies that patients with UC have an altered bacterial microbiota<sup>2)</sup>. Thus, the bacterial and/or host-bacterial interactions may play role in the pathogenesis of UC. This study focus on the metabolic interactions, corresponding to the fecal microbiota derived from UC patients and healthy subjects, colonizing a dynamic *in vitro* gut model.

## Materials and methods

Fecal samples came from 4 healthy volunteers and 8 UC patients (4 in remission and 4 in relapse state). Studies have been done in a dynamic *in vitro* gastrointestinal model, the M-SHIME<sup>2),3)</sup>. For metabolic analyses mucus and lumen samples were taken from the M-SHIME after 42 hours. In order to extract metabolites cold MeOH was used. Metabolites were detected by LCMS as follow: a Dionex Ultimate 3000 RS liquid chromatograph coupled to a Bruker maXis time of flight mass spectrometer. Analytes were separated on a Kinetex PFP column 50 x 2.10 mm, 2.6 µm, 100Å, using solvents: 10 mM NH<sub>4</sub>HCO<sub>2</sub> and C<sub>2</sub>H<sub>3</sub>N as a linear gradient from 0 to 90% C<sub>2</sub>H<sub>3</sub>N over 8 min. Scan range was from 50 to 800 m/z. The differences in metabolite profiles<sup>4)</sup> were evaluated by principal component analysis (PCA) using Profile Analysis 2.0 by Bruker Daltonics.

## Results and discussion

PCA showed a distinctive separation between UC in relapse and healthy donors, which confirmed the data, describing bacterial differences between two types of microflora, made for the same samples. The same result was observed for the lumen samples. Metabolites separating those two groups are e.g. bile acids, fatty acids and tryptophan. These results will be further studied and combined with the qPCR data, describing the changes in the bacterial community for the healthy donors and UC patients.

## Acknowledgments

This work was supported by the Danish Council for Strategic Research, Technical University of Denmark, Øresund Food and a GOA project from Ghent University.

## References

**1)** Kornbluth A, Sachar DB. (2010) Ulcerative Colitis Practice Guidelines in Adults: American College of Gastroenterology, Practice Parameters Committee. *American Journal of Gastroenterology*. 105:501-523. **2)** Vigsnaes LK. (2011) Role of Intestinal Microbiota in Ulcerative Colitis – Effects of Novel Carbohydrate Preparations. Ph.D. Thesis. Technical University of Denmark, DTU Food. **3)** Van den Abbeele P, Roos S *et al.*. (2011) Incorporating a mucosal environment in a dynamic gut model results in a more representative colonization by lactobacilli. *Microbial Biotechnology*. DOI: 10.1111/j.1751-7915.2011.00308.x. **4)** Sulek K, Frandsen HL *et al.*. (2011) Metabolic footprint of *Lactobacillus acidophilus* NCFM at different pH. *Metabolomics*. 8:244-252